

Allocasuarina mackliniana subsp. *hirtilinea* Western Sheoak

Taxonomy

Allocasuarina mackliniana subsp. *hirtilinea* L.A.S. Johnson

This is one of three subspecies of *A. mackliniana*, two of which occur in Victoria; one is shared with South Australia, and one endemic in SA.

Current conservation status

Categorised as Rare in the 2014 Advisory list of rare or threatened flora (DEPI 2014).

Proposed conservation status

Endangered in Australia

Criteria B1ab(i,ii,iii,iv)+2ab(i,ii,iii,iv)

Species Information

Description and Life History

The taxon is a long-lived dioecious, rarely monoecious, shrub to 3 m high, resprouting after fire. The taxon reproduces by seed only from a canopy-stored seedbank with massed recruitment only following fire. It is an obligatory outbreeder (dioecious) and it is pollinated by wind, with gene flow occurring via pollen and wind-dispersed seed, which potentially occurs over some distance.

Generation Length

The generation length of *Allocasuarina mackliniana* subsp. *hirtilinea* is suspected to be 50 to 100 years. This is based on the taxon's longevity, and episodic recruitment in response to fire.

Distribution

The taxon is confined to the Grampians with an outlying population in the southern Wimmera east of Douglas.

Habitat

The taxon exists in heathy and grassy woodland environments on sandy soils derived from sandstone or alluvial sediments.

Threats

Threats to the taxon include climate change (i.e. decreased rainfall, increased evaporation, extreme temperatures), altered fire regimes (i.e. increased frequency and intensity), inappropriate timing of prescribed fire (i.e. winter-spring), the impacts of fire-control activities, soil loss on bare post-fire substrates, weed invasion, and the grazing of seedlings by rabbits and hares.

Allocasuarina mackliniana subsp. *hirtilinea* Western Sheoak

IUCN Criteria

Criterion A. Population size reduction. Population reduction (measured over the longer of 10 years or 3 generations) based on any of A1 to A4			
	Critically Endangered	Endangered	Vulnerable
A1	≥ 90%	≥ 70%	≥ 50%
A2, A3, A4	≥ 80%	≥ 50%	≥ 30%
<p>A1 Population reduction observed, estimated, inferred or suspected in the past and the causes of the reduction are clearly reversible AND understood AND ceased.</p> <p>A2 Population reduction observed, estimated, inferred or suspected in the past where the causes of the reduction may not have ceased OR may not be understood OR may not be reversible.</p> <p>A3 Population reduction, projected or suspected to be met in the future (up to a maximum of 100 years) [(a) cannot be used for A3]</p> <p>A4 An observed, estimated, inferred, projected or suspected population reduction where the time period must include both the past and the future (up to a max. of 100 years in future), and where the causes of reduction may not have ceased OR may not be understood OR may not be reversible.</p>			
<p>based on any of the following:</p> <ul style="list-style-type: none"> (a) direct observation [except A3] (b) an index of abundance appropriate to the taxon (c) a decline in area of occupancy, extent of occurrence and/or quality of habitat (d) actual or potential levels of exploitation (e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites 			

Evidence:

Ineligible under Criterion A

The past population reduction does not meet the threshold for eligibility under criterion A2, and the future population reduction does not meet the threshold for eligibility under criterion A3.

Allocasuarina mackliniana subsp. *hirtilinea* Western Sheoak

Criterion B. Geographic range in the form of either B1 (extent of occurrence) and/or B2 (area of occupancy)			
	Critically Endangered Very restricted	Endangered Restricted	Vulnerable Limited
B1. Extent of occurrence (EOO)	< 100 km ²	< 5,000 km ²	< 20,000 km ²
B2. Area of occupancy (AOO)	< 10 km ²	< 500 km ²	< 2,000 km ²
AND at least 2 of the following 3 conditions:			
(a) Severely fragmented OR Number of locations	= 1	≤ 5	≤ 10
(b) Continuing decline observed, estimated, inferred or projected in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) area, extent and/or quality of habitat; (iv) number of locations or subpopulations; (v) number of mature individuals			
(c) Extreme fluctuations in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) number of locations or subpopulations; (iv) number of mature individuals			

Evidence:

Eligible under Criterion B1 as Endangered

The Extent of Occurrence (EoO) across the taxon's range is estimated to be 1,557 km², based on accepted, post-1970 records from the Victorian Biodiversity Atlas (VBA).

The taxon is inferred to be severely fragmented, and is projected to have 1 location. It has a continuing decline in (i), (ii), (iii) and (iv) above due to the identified threats, namely, climate change, altered fire regimes and weed invasion.

There is no capacity for long-distance dispersal and therefore no realistic capacity for recolonisation in the event of local extinction.

Eligible under Criterion B2 as Endangered

The Area of Occupancy (AoO) across the taxon's range is estimated to be 40 km², based on 2 x 2 km grids derived from accepted, post-1970 records in the VBA. As above, the taxon is inferred to be severely fragmented, has 1 location and has a continuing decline in (i), (ii), (iii) and (iv) above.

Allocasuarina mackliniana subsp. hirtilinea

Western Sheoak

Criterion C. Small Population size and decline		Critically Endangered	Endangered	Vulnerable
Number of mature individuals		< 250	< 2,500	< 10,000
AND at least one of C1 or C2				
C1	An observed, estimated or projected continuing decline of at least (up to a max. of 100 years in future):	25% in 3 years or 1 generation (whichever is longer)	20% in 5 years or 2 generations (whichever is longer)	10% in 10 years or 3 generations (whichever is longer)
C2	An observed, estimated, projected or inferred continuing decline AND least 1 of the following 3 conditions:			
(a)	(i) Number of mature individuals in each subpopulation	≤ 50	≤ 250	≤ 1,000
	(ii) % of mature individuals in one subpopulation =	90 – 100%	95 – 100%	100%
(b)	Extreme fluctuations in the number of mature individuals			

Evidence:

Ineligible under Criterion C as Data Deficient

No reliable estimate of the total population size for the taxon is available.

Criterion D. Very small or restricted populations		Critically Endangered	Endangered	Vulnerable
Number of mature individuals (observed or estimated)		< 50	< 250	< 1,000
D2. Only applies to the VU category Restricted area of occupancy or number of locations with a plausible future threat that could drive the species to critically endangered or Extinct in a very short time.		-	-	D2. Typically: AoO < 20 km ² or number of locations ≤ 5

Evidence:

Ineligible under Criterion D

There is insufficient evidence to determine the number of mature individuals.

Criterion E (Quantitative Analysis) was not addressed as the taxon does not have a detailed Population Viability Analysis.

References

DEPI (2014). *Advisory list of rare or threatened plants in Victoria - 2014*. Department of Environment and Primary Industries, Melbourne. Retrieved from: https://www.environment.vic.gov.au/__data/assets/pdf_file/0021/50448/Advisory-List-of-Rare-or-Threatened-Plants-in-Victoria-2014.pdf



Allocasuarina mackliniana subsp. *hirtilinea*
Western Sheoak

VicFlora (2019). Flora of Victoria, Royal Botanic Gardens Victoria: *Allocasuarina mackliniana* subsp. *hirtilinea*. Retrieved from: <https://vicflora.rbg.vic.gov.au/flora/taxon/be5f66b0-15e9-4554-af5b-4d9c576acef5>

Wilson, K.L., & Johnson, L.A.S. in George, A.S. (ed.) (1989), Casuarinaceae. *Flora of Australia* 3, 162.