

Pimelea ligustrina subsp. *ciliata* Fringed Rice-flower

Taxonomy

Pimelea ligustrina subsp. *ciliata* Threlfall

Two of the three subspecies occur in Victoria

Current conservation status

Categorised as Rare in the 2014 Advisory list of rare or threatened flora (DEPI 2014).

Proposed conservation status

Endangered in Victoria

Criteria B1ab(iii,v)+2ab(iii,v)

Species Information

Description and Life History

Pimelea ligustrina subsp. *ciliata* is a long-lived shrub to 1.5 m high, fire sensitive and recovering from a soil-stored seedbank of unknown longevity (but probably long-lived). Plants monoecious but flowers in inflorescence either bisexual or female. Breeding system unknown, i.e. whether self-fertile or an obligate outcrosser. Seeds (fruits) are passively dispersed but are likely to be wind-dispersed locally. Flowers are pollinated by Lepidoptera (mostly moths) visiting for a nectar reward.

Generation Length

The generation length of *Pimelea ligustrina* subsp. *ciliata* is suspected to be 30 to 50 years, based on the recurrent fire interval triggering seeding.

Distribution

The taxon occurs in alpine and sub-alpine areas of north-east Victoria.

Habitat

The taxon occurs in mesic alpine and subalpine Snow Gum and Alpine Ash woodland and forest, from c. 1400-2000 m above sea level, with shrubby-grassy understories on a range of well-drained loamy soils derived from a variety of geologies. It extends to just above the tree line into heath vegetation.

Threats

Threats include climate change (decreased rainfall and snow-fall, increased evaporation, and extreme temperatures); increased frequency and intensity of fire; soil loss on bare post-fire substrates resulting from severe rainfall events; damage by feral deer, cattle and horses; fungal pathogens; and weed invasion.

Pimelea ligustrina subsp. ciliata Fringed Rice-flower

IUCN Criteria

Criterion A. Population size reduction. Population reduction (measured over the longer of 10 years or 3 generations) based on any of A1 to A4			
	Critically Endangered	Endangered	Vulnerable
A1	≥ 90%	≥ 70%	≥ 50%
A2, A3, A4	≥ 80%	≥ 50%	≥ 30%

<p>A1 Population reduction observed, estimated, inferred or suspected in the past and the causes of the reduction are clearly reversible AND understood AND ceased.</p> <p>A2 Population reduction observed, estimated, inferred or suspected in the past where the causes of the reduction may not have ceased OR may not be understood OR may not be reversible.</p> <p>A3 Population reduction, projected or suspected to be met in the future (up to a maximum of 100 years) [(a) cannot be used for A3]</p> <p>A4 An observed, estimated, inferred, projected or suspected population reduction where the time period must include both the past and the future (up to a max. of 100 years in future), and where the causes of reduction may not have ceased OR may not be understood OR may not be reversible.</p>	<p>based on any of the following:</p>	<p>(a) direct observation [except A3]</p> <p>(b) an index of abundance appropriate to the taxon</p> <p>(c) a decline in area of occupancy, extent of occurrence and/or quality of habitat</p> <p>(d) actual or potential levels of exploitation</p> <p>(e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites</p>
--	---------------------------------------	---

Evidence:

Ineligible under Criterion A

There is insufficient evidence to determine whether there has been or will be a reduction in population sufficient to meet any threshold for Criterion A.

Pimelea ligustrina subsp. ciliata Fringed Rice-flower

Criterion B. Geographic range in the form of either B1 (extent of occurrence) and/or B2 (area of occupancy)			
	Critically Endangered Very restricted	Endangered Restricted	Vulnerable Limited
B1. Extent of occurrence (EOO)	< 100 km ²	< 5,000 km ²	< 20,000 km ²
B2. Area of occupancy (AOO)	< 10 km ²	< 500 km ²	< 2,000 km ²
AND at least 2 of the following 3 conditions:			
(a) Severely fragmented OR Number of locations	= 1	≤ 5	≤ 10
(b) Continuing decline observed, estimated, inferred or projected in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) area, extent and/or quality of habitat; (iv) number of locations or subpopulations; (v) number of mature individuals			
(c) Extreme fluctuations in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) number of locations or subpopulations; (iv) number of mature individuals			

Evidence:

Eligible under Criterion B1 as Endangered

The Extent of Occurrence (EoO) across the taxon's range is estimated to be 4,323 km², based on accepted, post-1970 records from the Victorian Biodiversity Atlas (VBA).

The taxon is suspected to be severely fragmented, as there is no capacity for long-distance dispersal and therefore no realistic capacity for recolonisation in the event of local extinction.

It is inferred to have a continuing decline in (iii) and (v) above, based on the impacts of the identified threats.

Eligible under Criterion B2 as Endangered

The Area of Occupancy (AoO) across the taxon's range is estimated to be 108 km², based on 2 x 2 km grids derived from accepted, post-1970 records in the VBA. As above, it severely fragmented, and has a continuing decline in (iii) and (v) above.

Pimelea ligustrina subsp. ciliata

Fringed Rice-flower

Criterion C. Small Population size and decline				
		Critically Endangered	Endangered	Vulnerable
Number of mature individuals		< 250	< 2,500	< 10,000
AND at least one of C1 or C2				
C1	An observed, estimated or projected continuing decline of at least (up to a max. of 100 years in future):	25% in 3 years or 1 generation (whichever is longer)	20% in 5 years or 2 generations (whichever is longer)	10% in 10 years or 3 generations (whichever is longer)
C2	An observed, estimated, projected or inferred continuing decline AND least 1 of the following 3 conditions:			
(a)	(i) Number of mature individuals in each subpopulation	≤ 50	≤ 250	≤ 1,000
	(ii) % of mature individuals in one subpopulation =	90 – 100%	95 – 100%	100%
(b)	Extreme fluctuations in the number of mature individuals			

Evidence:

Ineligible under Criterion C as Data Deficient

There is no population information as there have been no dedicated surveys for this taxon.

Criterion D. Very small or restricted populations				
		Critically Endangered	Endangered	Vulnerable
Number of mature individuals (observed or estimated)		< 50	< 250	< 1,000
D2. Only applies to the VU category Restricted area of occupancy or number of locations with a plausible future threat that could drive the species to critically endangered or Extinct in a very short time.		-	-	D2. Typically: AoO < 20 km ² or number of locations ≤ 5

Evidence:

Ineligible under criterion D

There is no population information as there have been no dedicated surveys for this taxon.

Criterion E (Quantitative Analysis) was not addressed as the taxon does not have a detailed Population Viability Analysis.

References

Costin, A.B., Gray, M., Totterdell, C.J., and Wimbush, D.J. (1979). *Kosciusko Alpine Flora*. Melbourne: CSIRO/Collins Australia.

DEPI (2014) *Advisory list of rare or threatened plants in Victoria - 2014*. Department of Environment and Primary Industries, Melbourne.

McMahon, A. (pers. comm.)



Pimelea ligustrina subsp. *ciliata*
Fringed Rice-flower

Rye, B.L. (1990). *Pimelea*. In D. Chapman (Ed) *Flora of Australia* Vol 18 (pp. 134-211). Melbourne, CSIRO.

Threlfall, S. (1983). The *genus* *Pimelea* (Thymelacaceae) in eastern mainland Australia. *Brunonia* 5(2), 113-202.

VicFlora (2019) Flora of Victoria, Royal Botanic Gardens Victoria: *Pimelea ligustrina* subsp. *ciliate*. Retrieved from <https://vicflora.rbg.vic.gov.au/flora/taxon/5dbfa34f-e5a6-4d67-b3b1-722e7c12bd39>