



## *Thelymitra lucida* Glistening Sun-orchid

### Taxonomy

*Thelymitra lucida* Jeanes

The taxon is distinguished from *Thelymitra holmesii* by the sparkling bloom on the mid-lobe of the column and the less shaggy hair tufts on the lateral lobes (VicFlora, 2018).

### Current conservation status

Categorised as Endangered in the 2014 Advisory list of rare or threatened flora (DEPI 2014).

### Proposed conservation status

Endangered in Victoria

Criterion A2bc+3c+4bc; B2ab(iii,v); C2a(i)

### Species Information

#### Description and Life History

The taxon has an erect flowering stem, straight, 30-55 cm tall, 1-3.5 mm diam., green to purplish. Leaf linear to linear-lanceolate, attenuate, 20-35 cm long, 5-12 mm wide, fleshy, canaliculate, ribbed abaxially, sheathing at base, dark green with a purplish base. Inflorescence 1-7-flowered, open. Sterile bracts usually 2, sometimes 3. Perianth segments lanceolate to ovate, 8-12 mm long, dark blue. Column slender, 5-6 mm long, blue to pinkish; mid-lobe expanded into hood over the anther, tubular, compressed dorsally, gently curved, yellow with a dark purplish black collar, covered with a thin glistening bloom, apex deeply bilobed; lateral lobes converging 1-1.5 mm long, digitiform, porrect, more or less straight, each with a toothbrush-like arrangement of white or creamy yellow hairs terminating in front of mid-lobe. Anther inserted about mid-way along column, shortly beaked. The taxon flowers in November and December. Flowers are self-pollinating and only open on warm, humid days. This taxon is facultatively autogamous (VicFlora, 2018).

Although recognised as distinct for several years now, the taxon was only recently described. Plants will grow submerged for part of the year, and flowering often occurs while the plants are still in standing water. It is a poorly known orchid that appears to be extremely rare, having been recorded from very few sites and plants. However, the taxon can be extremely difficult to detect in its dense sedge swamp habitat when not in flower. (Backhouse et al., 2016).

#### Generation Length

The generation length of *Thelymitra lucida* is suspected to be 20 to 40 (midpoint 30) years. Generation time for non-colonial terrestrial orchids is estimated to be a nominal 30 years based on the annual replacement of the mother tuber by daughter tubers. Whilst somatically immortal, each individual is susceptible to endogenous exhaustion or environmental causes of mortality at rates likely to result in replacement at intervals of several decades only. Such orchids are classed as obligate seed regenerators (OSRs) reliant on seed-based recruitment for population maintenance.

# *Thelymitra lucida* Glistening Sun-orchid

## Distribution

The taxon occurs in western Victoria, mostly west from the Grampians, also in the Brisbane Ranges and on the basalt plains at Caramut. The altitude ranges from 50-370 metres above sea level. The taxon also occurs in Tasmania. The taxon is probably under-reported, and may be more commonly currently recorded, especially considering there is a record from a boggy site in herb-rich plains grasslands (Backhouse et al., 2016). The taxon was reasonably widespread and represented in reserves but extremely rare and seldom collected (Jeanes, 2004).

## Habitat

The taxon grows in or near lowland sedge swamps on peaty soil, mostly in standing water at flowering time, rarely in wet depressions in plains grassland (Backhouse et al., 2016; VicFlora, 2018).

## Threats

The taxon grows around the margins of shallow freshwater marshes, and there has been a widespread historical loss of this habitat across the range of the taxon, which is likely to have led to the loss of subpopulations.

There is the possibility of a future decline in distribution and abundance of the taxon, based on the likely long-term decline and loss of shallow freshwater marsh habitat from increasingly dry conditions due to declining rainfall.

## IUCN Criteria

Criterion A. Population size reduction. Population reduction (measured over the longer of 10 years or 3 generations) based on any of A1 to A4			
	Critically Endangered	Endangered	Vulnerable
A1	≥ 90%	≥ 70%	≥ 50%
A2, A3, A4	≥ 80%	≥ 50%	≥ 30%
<p>A1 Population reduction observed, estimated, inferred or suspected in the past and the causes of the reduction are clearly reversible AND understood AND ceased.</p> <p>A2 Population reduction observed, estimated, inferred or suspected in the past where the causes of the reduction may not have ceased OR may not be understood OR may not be reversible.</p> <p>A3 Population reduction, projected or suspected to be met in the future (up to a maximum of 100 years) [(a) cannot be used for A3]</p> <p>A4 An observed, estimated, inferred, projected or suspected population reduction where the time period must include both the past and the future (up to a max. of 100 years in future), and where the causes of reduction may not have ceased OR may not be understood OR may not be reversible.</p> <p>based on any of the following:</p> <ul style="list-style-type: none"> <li>(a) direct observation [except A3]</li> <li>(b) an index of abundance appropriate to the taxon</li> <li>(c) a decline in area of occupancy, extent of occurrence and/or quality of habitat</li> <li>(d) actual or potential levels of exploitation</li> <li>(e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites</li> </ul>			

## Evidence:

### Eligible under Criterion A2 as Endangered

The population reduction over the past 60 to 120 years is inferred to be 30 to 50%, based on (b) and (c) above. Past decline is based on the historical loss of shallow freshwater marshes across the range of the taxon.

# *Thelymitra lucida* Glistening Sun-orchid

## Eligible under Criterion A3 as Endangered

The population reduction over the next 60 to 100 years is projected to be 40 to 60%, based on (c) above.

Future decline is based on the likely long-term decline and loss of shallow freshwater marshes from increasingly dry conditions due to declining rainfall.

## Eligible under Criterion A4 as Endangered

The population reduction over any 60 to 120 year period, including both past and future (up to 100 years in the future), is inferred to be 40 to 60%, based on b) and (c) above.

Criterion B. Geographic range in the form of either B1 (extent of occurrence) and/or B2 (area of occupancy)			
	Critically Endangered Very restricted	Endangered Restricted	Vulnerable Limited
B1. Extent of occurrence (EOO)	< 100 km <sup>2</sup>	< 5,000 km <sup>2</sup>	< 20,000 km <sup>2</sup>
B2. Area of occupancy (AOO)	< 10 km <sup>2</sup>	< 500 km <sup>2</sup>	< 2,000 km <sup>2</sup>
AND at least 2 of the following 3 conditions:			
(a) Severely fragmented OR Number of locations	= 1	≤ 5	≤ 10
(b) Continuing decline observed, estimated, inferred or projected in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) area, extent and/or quality of habitat; (iv) number of locations or subpopulations; (v) number of mature individuals			
(c) Extreme fluctuations in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) number of locations or subpopulations; (iv) number of mature individuals			

## Evidence:

### Eligible under Criterion B as Endangered

The Area of Occupancy (AoO) is estimated to be 16 km<sup>2</sup>, based on 2 x 2 km grids derived from accepted, post-1970 records in the Victorian Biodiversity Atlas (VBA).

Any two of (a), (b) or (c) above are also satisfied.

# Thelymitra lucida

## Glistening Sun-orchid

Criterion C. Small Population size and decline				
		Critically Endangered	Endangered	Vulnerable
Number of mature individuals		< 250	< 2,500	< 10,000
AND at least one of C1 or C2				
C1	An observed, estimated or projected continuing decline of at least (up to a max. of 100 years in future):	25% in 3 years or 1 generation (whichever is longer)	20% in 5 years or 2 generations (whichever is longer)	10% in 10 years or 3 generations (whichever is longer)
C2	An observed, estimated, projected or inferred continuing decline AND least 1 of the following 3 conditions:			
(a)	(i) Number of mature individuals in each subpopulation	≤ 50	≤ 250	≤ 1,000
	(ii) % of mature individuals in one subpopulation =	90 – 100%	95 – 100%	100%
(b)	Extreme fluctuations in the number of mature individuals			

### Evidence:

#### Eligible under Criterion C2 as Endangered

It is estimated that there are 75 to 220 mature individuals, based on sporadic surveys and VBA records.

The number of mature individuals is inferred to continue to decline, and the number of mature individuals in each subpopulation is fewer than 250.

Criterion D. Very small or restricted populations				
		Critically Endangered	Endangered	Vulnerable
Number of mature individuals (observed or estimated)		< 50	< 250	< 1,000
D2. Only applies to the VU category Restricted area of occupancy or number of locations with a plausible future threat that could drive the species to critically endangered or Extinct in a very short time.		-	-	D2. Typically: AoO < 20 km <sup>2</sup> or number of locations ≤ 5

### Evidence:

#### Eligible under Criterion D as Endangered

It is estimated that there are 75 to 220 mature individuals.

Criterion E (Quantitative Analysis) was not addressed as the taxon does not have a detailed Population Viability Analysis.

### References

Backhouse, G., Kosky, B., Rouse, D., and Turner, J. (2016). *Bush Gems: A Guide to the Wild Orchids of Victoria, Australia*. Melbourne, Victoria: EBook.



# *Thelymitra lucida* Glistening Sun-orchid

DEPI (2014). *Advisory list of rare or threatened plants in Victoria - 2014*. Department of Environment and Primary Industries, Melbourne.

Jeanes, J. A. (2004). A revision of the *Thelymitra pauciflora* R.Br. (Orchidaceae) complex in Australia. *Muelleria*, 19, 19-79.

VicFlora (2018). Flora of Victoria, Royal Botanic Gardens Victoria: *Thelymitra lucida*. Retrieved from: <https://vicflora.rbg.vic.gov.au/flora/taxon/dc287237-a0c6-42f5-9867-99d73b1feb27>